



## Isolation of soil *Streptomyces* as source antibiotics active against antibiotic-resistant bacteria

Ozgur Ceylan\*, Gulden Okmen, Aysel Ugur

Department of Biology, Faculty of Arts and Sciences, Mugla University 48187  
Kotekli-Mugla, Turkey

\*Corresponding author: ozgceylan@hotmail.com

### Abstract

The focus of this study was the in vitro antimicrobial activities of Streptomyces, bacteria commonly found in soil and known antibiotic-producers. *Streptomyces* isolates obtained from different fields in Muğla, Turkey were evaluated for their inhibitory activities on seven microorganisms including multiple antibiotic resistant *Staphylococcus aureus* and *Stenotrophomonas maltophilia*. Fifteen *Streptomyces* isolates which exhibited antimicrobial activity against at least two of the test organisms were characterized by conventional methods. The results indicated that five isolates were highly active against *S. aureus* strains including methicillin resistant *Staphylococcus aureus* (MRSA). Twelve Streptomyces isolates showed anticandidal activity against *Candida albicans*. Ten isolates were highly active with an inhibition zone more than 30 mm in diameter. Most of the isolates inhibited growth of the Gram negative bacteria tested. Eight isolates showed antibacterial activity on *S. maltophilia* MU64. The inhibition zones of two were higher than 30 mm for *S. maltophilia*.

**Keywords:** Antimicrobial activity, microorganism, *Streptomyces*.

Ceylan O, Okmen G, Ugur A (2008) Isolation of soil *Streptomyces* as source antibiotics active against antibiotic-resistant bacteria. EurAsia J BioSci 2, 9, 73-82.  
[www.ejobios.com/content/2/9/73-82](http://www.ejobios.com/content/2/9/73-82)

### INTRODUCTION

Serious infections caused by bacteria that have become resistant to commonly used antibiotics has become a major global healthcare problem in the 21<sup>st</sup> century (Alanis 2005). *Staphylococcus aureus*, for instance, a virulent pathogen that is responsible for a wide range of infections including pimples, pneumonia, osteomyelitis, endocarditis and bacteremia, has developed resistance to most classes of antibiotics (Enright 2003). For more than two decades, clinicians and public health officials have faced hospital acquired methicillin-resistant *S. aureus* (MRSA), which also bears resistance to many antibiotics. During much of this time, vancomycin has been the therapeutic answer to MRSA, but that paradigm has changed. Vancomycin-resistant strains have emerged clinically (Hiramatsu 1998, Bozdogan et al. 2003, Chang et al. 2003, Anonymous 2004). Vancomycin-resistant *S. aureus* (VRSA) challenges clinicians, not only because of

vancomycin and methicillin resistance, but also because of resistance to many other antibiotics, including aminoglycosides, macrolides, and fluoroquinolones. Fortunately, newer therapeutic agents, daptomycin, linezolid, and a streptogramin combination (quinupristin/dalfopristin) have entered the clinical arena in the past few years (Levy and Marshall 2004, Wenzel 2004). However, certain undesirable side effects and the spread of pathogens with this new antimicrobial drug resistance emphasize the need for the development of other newer antimicrobial agents with activity against such Gram positive bacteria (Jevitt et al. 2003, Meka and Gold 2004, Wenzel 2004, Nathwani 2005). Another cause of great concern is the Gram negative antibiotic-resistant opportunistic pathogens. Gram negative environmental and enteric organisms currently threaten patients

Received: June, 2008  
Accepted: July, 2008  
Printed: September, 2008

in hospitals and communities with multi-drug resistance, including broad resistance to first, second, and third generations of penicillin's and cephalosporin's (Urban et al. 2003, Obritsch et al. 2004, Paterson et al. 2004). These bacteria, like *Pseudomonas aeruginosa*, are common environmental organisms, which act as opportunistic pathogens in clinical cases where the defense system of the patient is compromised (Lyczak et al. 2000). In addition, other intrinsically antibiotic resistant organisms such as *Stenotrophomonas maltophilia* (Saiman et al. 2002) are emerging as opportunistic pathogens.

The end result of this phenomenon is that many strains of bacteria have become resistant, and in many cases multi-resistant to these therapeutic agents, thus rendering these drugs ineffective as treatments of choice for severe infections caused by these pathogens (Alanis 2005). Rising numbers of antibiotic-unresponsive infectious disease agents confront patients worldwide (Levy 2002, Livermore 2003) and consensus has emerged that it is essential that novel antibiotic classes be developed as part of the strategy to control the emerging drug-resistant pathogens (Projan 2002, Abbanat et al. 2003, Barrett and Barrett 2003). In response, there is a renewed interest in discovering novel classes of antibiotics that have different mechanisms of action (Spizek and Tichy 1995, Barsby et al. 2001).

Search for new antibiotics effective against multi-drug resistant pathogenic bacteria is presently an important area of antibiotic research. Natural products having novel structures have been observed to possess useful biological activities. Soil is a natural reservoir for microorganisms and their antimicrobial products (Dancer 2004). Filamentous soil bacteria belonging to the genus *Streptomyces* are widely recognized as industrially important microorganisms because of their ability to produce many kinds of novel secondary metabolites including antibiotics (Williams et al. 1983a, Crandall and Hamil 1986, Williams et al. 1989, Korn-Wendisch and Kutzner 1992). Of all known drugs 70% have been isolated from Actinomycetes

bacteria of which 75% and 60% are used in medicine and agriculture respectively (Miyadoh 1993, Tanaka and Mura 1993).

The genus *Streptomyces* was proposed by Waksman and Henrici for aerobic and spore-forming Actinomycetes (Williams et al. 1989). The taxon currently accommodates Gram positive bacteria that have a DNA with a high guanine-plus-cytosine content (69 to 73 mol %) and that form extensive branching substrates and aerial mycelia (Williams et al. 1983a, Williams et al. 1989, Korn-Wendisch and Kutzner 1992).

Indeed, different *Streptomyces* species produce about 75% of commercially and medically useful antibiotics. They have provided more than half of the naturally occurring antibiotics discovered to date and continue to be screened for useful compounds (Miyadoh 1993). In the course of screening for new antibiotics, several studies are oriented towards isolation of Streptomyces from different habitats. Presently, there is little documented information of the occurrence of *Streptomyces* spp. in the soil of Turkey with a potential to produce antimicrobial compounds (Denizci 1996, Aslan 1999, Sahin and Ugur 2003, Oskay et al. 2004). In the present study, the isolation and characterization as well as the inhibitory effects of local Streptomyces isolates tested against various multiple antibiotic resistant bacteria and yeast were reported, along with some chemical properties of secondary metabolites with high biological activities.

---

## MATERIAL AND METHODS

---

### Isolation of microorganisms

Soil samples were collected from various locations in Mugla province from 2006 to 2007. Several diverse habitats in different areas were selected for the isolation of *Streptomyces* strains. These habitats included the rhizosphere of plants, agricultural soil, preserved areas and forest soils. The samples were taken up to a depth of 20 cm after removing approximately 3 cm of the soil surface. The samples were placed in polyethylene bags, closed tightly and stored in

a refrigerator. The following screening procedure was adopted for the isolation of *Streptomyces* (Korn-Wendisch and Kutzner 1992). The soil was pretreated with CaCO<sub>3</sub> (10:1 w/w) and incubated at 37°C for 4 days. It was then suspended in sterile Ringer solution (1/4 strength). Test tubes containing a 10-2 dilution of the samples were placed in a water bath at 45°C for 16 h so that the spores would separate from the vegetative cells and the dilutions were inoculated on the surface of the Actinomycete Isolation Agar (Difco 0957) plates. The plates were incubated at 28°C until the sporulation of *Streptomyces* colonies occurred. *Streptomyces* colonies (where the mycelium remained intact and the aerial mycelium and long spore chains were abundant) were then removed and transferred to the Yeast Extract-Malt Extract Agar (ISP2) slants. Pure cultures were obtained from selected colonies for repeated sub culturing. After antimicrobial activity screening, the isolated *Streptomyces* strains were maintained as suspensions of spores and mycelial fragments in 10% glycerol (v/v) at -20°C in the Mugla University Collection of Microorganisms (MU).

#### Characterization of the isolates

*Streptomyces* colonies were characterized morphologically and physiologically following the directions given for the International *Streptomyces* Project (ISP) (Shirling and Gottlieb 1966). General morphology was determined using the Oatmeal Agar plates, incubated in the dark at 28°C for 21 days and then by direct light microscopy examination of the surface of the cross-hatched cultures. Colours were determined according to the scale adopted by Prauser (1964). Melanin reactions were detected by growing the isolates on Peptone-Yeast Extract-Iron Agar (ISP 6) (Shirling and Gottlieb 1966). All strains were cultivated on an ISP 2 medium. Some diagnostic characters of highly active *Streptomyces* strains were determined following the directions given in the Bergey's Manual of Systematic Bacteriology (Williams et al. 1983a, 1983b).

#### Test microorganisms

Six bacteria, including three Gram positive

(*Staphylococcus aureus* MU 38, *Staphylococcus aureus* MU 40, *Staphylococcus aureus* ATCC 25923) and three Gram negative (*Escherichia coli* ATCC 25922, *Pseudomonas aeruginosa* ATCC 27853, *Stenotrophomonas maltophilia* MU 64) and one yeast (*Candida albicans* ATCC 1023) were used to determine the antimicrobial activity of the isolated *Streptomyces* strains. All these microorganisms were obtained from the American Type Culture Collection (ATCC) and the Mugla University Collection of Microorganisms (MU) in Mugla, Turkey. The above mentioned bacteria were cultured in a Nutrient Broth (NB) (Difco) at 37±0.1°C for 24 h with *C. albicans* being cultured in a Sabouraud Dextrose Broth (SDB) (Difco) at 28±0.1°C for 48 h.

#### In vitro screening of isolates for antagonism

Balanced sensitivity medium (BSM, Difco 1863) plates were prepared and inoculated with *Streptomyces* isolate by a single streak of inoculum in the center of the petri dish. After 4 days of incubation at 28°C the plates were seeded with test organisms by a single streak at a 90° angle to the *Streptomyces* strains. The microbial interactions were analyzed by the determination of the size of the inhibition zone (Madigan et al. 1997).

---

## RESULTS AND DISCUSSION

---

The *Streptomyces* flora of 11 soil samples, collected from different locations in the Mugla region, were screened for their potential as a source of antibiotics active against antibiotic-resistant bacteria. All of the isolates were tested for their ability to produce inhibitory substances against seven test microorganisms (Data not shown). The test microorganisms included 3 Gram positive bacteria, 3 Gram negative bacteria and 1 yeast. Of them *S. aureus* MU38, MU40 and *S. maltophilia* MU64, are resistant to the widely used antibiotics. The antibiotic resistance patterns of these strains are shown in Table 1. The isolates which exhibited antimicrobial activity against at least two of the test

organisms were selected for this study. As shown in Table 2, a total of 15 different Streptomyces isolates were shown to have a very potent in vitro antimicrobial activity against the test organisms. The morphological examination of these isolates, which were active on the test organisms, indicates that these belong to the *Streptomyces* genus (Waksman 1961, Shirling and Gottlieb 1966, Nonomura 1974, Williams et al. 1983a, Cross 1989, Goodfellow 1989, Lechevalier 1989, Locci 1989). The morphological and cultural characteristics of the *Streptomyces* isolates are shown in Table 3.

As shown in Table 3, the percentage of active isolates varies within each colour series, with production of such compounds being recorded as a soluble pigment in the colours of brown-yellow (80%), yellow (6.6%), and violet (13.3%). The rate of melanin pigment production was 13.3%. The colour of the aerial misellium was white (73.3%), grey (20%) and violet (6.6%). The reverse side colour was brown-yellow (93.3%) and violet (6.6%).

Anticandidal activities were exhibited by all of the isolates, except MS1, MS12 and MS15. Thirty isolates produced antibacterial substances against both Gram negative and Gram positive bacteria. Ten of them inhibited the growth of bacteria with  $\geq 30$  mm inhibition zones. *S. aureus* strains were inhibited by almost all of the isolates. The isolates produced 3-48 mm, 4- 56 mm and 7-40 mm inhibition zones on *S. aureus* ATCC 25923, MU38 and MU40, respectively. Two of the isolates (MS10 and MS13) were not active against all of the tested *S. aureus* strains. The isolates MS12, MS14 and MS15 did not show any inhibition effect on *S. aureus* ATCC 25923 and MU40. Other isolates were active against one or more of them. Especially, the isolates MS4, MS9 and MS11 inhibited the growth of all of the tested *S. aureus* strains with  $\geq 30$  mm inhibition zones. Data indicated that, the many characterization of the three isolates were different, except the reverse side colour and the ability to produce melanin pigments. In addition, the isolates MS7 and MS8 inhibited

**Table 1.** The antibiotic resistance patterns of *S. aureus* MU 38, MU 40 and *S. maltophilia* MU 64.

Microorganisms	Resistance patterns <sup>a</sup>
<i>S. aureus</i> MU 38	P, AK, DA, CN, ME, TEC, TE
<i>S. aureus</i> MU 40	P, AK, CN, KF, ME, TE, OX
<i>S. maltophilia</i> MU 64	MEZ, TIM, CAZ, FEP, CRO, CTX, KF, IPM, P, AK, TOB, NET, CN, TE, CIP, NOR, C, SXT, TVA, AM, PRL, ATM, SAM, AMC

P: Penicillin (10 U), AK: Amikacin (30 mcg), DA: Clindamycin (2 mcg), CN: Gentamicin (10 mcg), KF: Cephalothin (30 mcg), ME: Methicillin (5 mcg), TE: Tetracycline (30 mcg), OX: Oxacillin (1 mcg), TEC: Teicoplanin (30 mcg). MEZ: Mezlocillin (75 mcg), TIM: Ticarcillin+clavulanic acid (75+10 mcg), CAZ: Ceftazidime (30 mcg), FEP: Cefepim (30 mcg), CRO: Ceftriaxone (30 mcg), CTX: Cefotxime (30 mcg), IPM: Imipenem (10 mcg), TOB: Tobramycin (10 mcg), NET: Netilmicin (30 mcg), NOR: Norfloxacin (10 mcg), C: Chloramphenicol (30 mcg), TVA: Trovafloksasin (10 mcg), AM: Ampicillin (10 mcg), PRL: Piperacillin (100 mcg), ATM: Aztreonam (30 mcg), SAM: Sulbactam + Ampicillin (10+10 mcg), AMC: Amoxicillin + Clavulanic acid (20+10 mcg), CIP: Ciprofloxacin (5 mcg), SXT: Trimetoprim + sulfamethoxazole (1.25+23.75 mcg).

<sup>a</sup> The NCCLS numeric values for the inhibition zones (in mm) of the bacteria to the antibiotics are given above in square brackets (NCCLS, 1999). If the inhibition zones determined in this study are the same or smaller than the inhibition zones cited, then the strains are considered resistant to the antibiotic tested.

**Table 2.** Antimicrobial activity of *Streptomyces* isolates.

Isolate no.	Microorganisms							
	<i>S. aureus</i>			<i>E. coli</i>	<i>P. aeruginosa</i>	<i>S. maltophilia</i>	<i>C. albicans</i>	
	ATCC 25923	MU 38	MU 40	ATCC 25922	ATCC 27853	MU 64	ATCC 1023	
	Inhibition zone (mm)							
MS1	3	-	-	3	-	-	-	-
MS2	20	15	30	18	9	>30	>30	>30
MS3	11	12	18	11	-	-	-	>30
MS4	48	>30	40	45	28	20	21	21
MS5	12	11	7	10	8	-	-	>30
MS6	16	18	14	16	15	-	-	>30
MS7	35	28	54	12	18	-	-	>30
MS8	48	30	29	31	5	15	>30	>30
MS9	32	56	31	36	53	3	32	32
MS10	-	-	-	-	12	-	-	>30
MS11	39	34	>30	40	6	5	22	22
MS12	-	4	-	4	8	7	-	-
MS13	-	-	-	-	5	-	-	>30
MS14	-	6	-	-	4	10	>30	>30
MS15	-	8	-	-	28	48	-	-

**Table 3.** Characteristics of the active *Streptomyces* isolates.

Isolate no.	Aerial mycelium	Reverse side colour	Soluble pigment colour	Melanin pigment	Sporophore morphology
MS1	Grey	Brown-Yellow	Brown-Yellow	-	Retinaculaperti
MS2	White	Brown-Yellow	Brown-Yellow	-	Retinaculaperti
MS3	White	Brown-Yellow	Brown-Yellow	-	Retinaculaperti
MS4	White	Brown-Yellow	Brown-Yellow	-	Retinaculaperti
MS5	White	Brown-Yellow	Brown-Yellow	+	Retinaculaperti
MS6	Violet	Violet	Violet	-	Retinaculaperti
MS7	Grey	Brown-Yellow	Violet	-	Retinaculaperti
MS8	White	Brown-Yellow	Brown-Yellow	-	Retinaculaperti
MS9	Grey	Brown-Yellow	Yellow	-	Spirales
MS10	White	Brown-Yellow	Brown-Yellow	-	Retinaculaperti
MS11	White	Brown-Yellow	Brown-Yellow	-	Spirales
MS12	White	Brown-Yellow	Brown-Yellow	+	Retinaculaperti
MS13	White	Brown-Yellow	Brown-Yellow	-	Rectiflexibles
MS14	White	Brown-Yellow	Brown-Yellow	-	Retinaculaperti
MS15	White	Brown-Yellow	Brown-Yellow	-	Rectiflexibles

the growth of tested *S. aureus* strains with  $> 28$  mm inhibition zones. All of the five

isolates did not produced melanin pigment. The cultural and morphological properties of the isolates MS4 and MS8 were the same. It appears that the inhibitory substances from MS4, MS7, MS8, MS9 and MS11 are more effective against *S. aureus* strains. The inhibitory substances produced by these five isolates were highly potent; they inhibited the *S. aureus* MU38, MU40 and ATCC 25923. *S. aureus* is a major cause of nosocomial infections, food poisoning, osteomyelitis, pyoarthritis, endocarditis, toxic shock syndrome and a broad spectrum of other disorders (Willett 1992, Todd 1998, Hajjeh et al. 1999, Rubin et al. 1999). *S. aureus* MU38 and MU40 are Methicillin-Resistant *Staphylococcus aureus* (MRSA). MRSA is responsible for the largest outbreak of hospital-acquired infection (HAI) that the world has ever seen (Gould 2005). MRSA is probably the most popular hospital resistant bacteria (Wenzel 2004, Beovic 2006). In recent years, there has been an alarming increase in nosocomial staphylococcal infections by strains with multiple drug resistance (Lyon and Skurray 1987, Al-Masaudi et al. 1991, Kloos and Bannerman 1995, Hiramatsu et al. 1997). Worldwide, many strains of *S. aureus* are already resistant to all antibiotics and the organism has progressed one step toward becoming an unstoppable killer (Ugur and Ceylan 2003).

The isolates produced antibacterial substances against one or more Gram negative bacteria tested. The isolates produced 3- 45 mm, 4-53 mm and 3-48 mm inhibition zones on *E. coli* ATCC 25922, *P. aeruginosa* ATCC 27853 and *S. maltophilia* MU64 respectively. Six isolates inhibited the growth of all the tested Gram negative bacteria. Two isolates (MS1, MS3) were only effective on *E. coli* among the Gram negative test bacteria. Four isolates (MS4, MS8, MS9, and MS11) produced  $\geq 30$  mm inhibition

zones for *E. coli*. The highest inhibition zone (53 mm) was on *P. aeruginosa*. The isolates MS4 and MS15 inhibited the growth of *P. aeruginosa* with 28 mm inhibition zones. Eight isolates showed antibacterial activity on *S. maltophilia* MU64 which is a multiple antibiotic resistant bacteria. The inhibition zones of two of them (MS2, MS15) were higher than 30 mm for *S. maltophilia*. *S. maltophilia*, (Palleroni and Bradbury 1993) (Xanthomonadaceae) previously known as *Pseudomonas maltophilia* (Hugh and Ryschenkow 1961) (Pseudomonadaceae) and subsequently as *Xanthomonas maltophilia* (Swings et al. 1983) (Xanthomonadaceae) has received much attention in the last decade because of its role as a pathogenic microorganism in an increasing number of clinical syndromes (Robin and Janda 1996), such as bacteremia, infections of the respiratory and urinary tracts, skin and soft tissue infections, biliary tract infection, meningitis, serious wound infections, conjunctivitis, endocarditis (Fisher et al. 1981, Denton and Kerr 1998), cystic fibrosis and central nervous system infections. *S. maltophilia* has also been described to be an important nosocomial pathogen (Denton and Kerr 1998). The treatment of infections caused by this microorganism is difficult because *S. maltophilia* is frequently resistant to most of the widely used antibiotics (Vartivarian et al. 1994, Liu et al. 1995, Skaehill 2000, Krueger et al. 2001).

In this work, we have shown that a total of 15 different *Streptomyces* isolates associated with soil have the ability to produce antimicrobial compounds against microorganisms, especially multiple antibiotic resistant Gram positive and Gram negative bacteria. Further investigations are needed in order to further determine the active metabolites of these isolates.

---

## REFERENCES

---

- Abbanat D, Macielag M, Bush K (2003) Novel antibacterial agents for the treatment of serious Gram-positive infections. *Expert Opinion on Investigational Drugs* 12, 379-399.

- Alanis AJ (2005) Resistance to antibiotics: are we in the post-antibiotic era? *Archives of Medical Research* 36, 697-705.
- Al-Masaudi SB, Day MJ, Russell AD (1991) Antimicrobial resistance and gene transfer in *Staphylococcus aureus*. *Journal of Applied Bacteriology* 70, 270-290.
- Anonymous (2004) Brief report: vancomycin-resistant *Staphylococcus aureus*-New York. *MMWR* 53, 322-323.
- Aslan B (1999) Studies on isolation, characterization and antibiotic production of *Streptomyces* species. PhD thesis, Cukurova University, Institute of Science, Adana.
- Barrett CT, Barrett JF (2003) Antibacterials: are the new entries enough to deal with the emerging resistance problems? *Current Opinion in Biotechnology* 14, 621-626.
- Barsby T, Kelly MT, Gagne SM (2001) Bogorol A produced in culture by a marine *Bacillus* sp. reveals a novel template for cationic peptide antibiotics. *Organic Letters* 3, 3, 437-440.
- Beovic B (2006) The issue of antimicrobial resistance in human medicine. *International Journal of Food Microbiology* 112, 3, 280-287.
- Bozdogan B, Esel D, Whitener C (2003) Antibacterial susceptibility of a vancomycin-resistant *Staphylococcus aureus* strain isolated at the Hershey Medical Center. *Journal of Antimicrobial Chemotherapy* 52, 864-868.
- Chang S, Sievert DM, Hageman JC (2003) Infection with vancomycin resistant *Staphylococcus aureus* containing the vanA resistance gene. *The New England Journal of Medicine* 348, 1342-1347.
- Crandall LW, Hamil RL (1986) Antibiotics produced by *Streptomyces*: major structural classes. In: Queener SW, Day LE (eds), *The bacteria*, Vol. 9, Academic Press, Orlando, Fla, 355-401.
- Cross T (1989) Growth and examination of actinomycetes some guidelines. In: *Bergey's Manual of Systematic Bacteriology*, Vol. 4, Williams and Wilkins Company, Baltimore, 2340-2343.
- Dancer SJ (2004) How antibiotics can make us sick: the less obvious adverse effects of antimicrobial chemotherapy. *The Lancet Infectious Diseases* 4, 611-619.
- Denizci AA (1996) A study on the detection and production of antibacterial antibiotics from Actinomycetes which isolated from the soils of Aegean and Eastern Black Sea regions of Turkey. PhD thesis, Ege University, Institute of Science, Izmir.
- Denton M, Kerr KG (1998) Microbiological and clinical aspects of infection associated with *Stenotrophomonas maltophilia*. *Clinical Microbiology Reviews* 11, 57-80.
- Enright MC (2003) The evolution of a resistant pathogen-the case of MRSA. *Current Opinion in Pharmacology* 3, 5, 474-479.
- Fisher MC, Long SS, Roberts EM (1981) *Pseudomonas maltophilia* bacteremia in children undergoing open heart surgery. *The Journal of the American Medical Association* 246, 1471-1474.
- Goodfellow M (1989) Suprageneric classification of Actinomycetes. In: *Bergey's Manual of Systematic Bacteriology*, Vol. 4, Williams and Wilkins Company, Baltimore, 2333-2339.
- Gould IM (2005) The clinical significance of methicillin-resistant *Staphylococcus aureus*. *Journal of Hospital Infection* 61, 277-282.
- Hajjeh RA, Rheingold A, Weil A, Shutt K, Schuhat A, Perkins BA (1999) Toxic shock syndrome in the United States, 1979-1996. *Emerging Infectious Diseases* 5, 807-810.

- Hiramatsu K, Hanaki H, Ino T, Yabuta K, Oguri T, Tenover FC (1997) Methicillin-resistant *Staphylococcus aureus* clinical strain with reduced vancomycin susceptibility. *The Journal of Antimicrobial Chemotherapy* 40, 135-136.
- Hiramatsu K (1998) Vancomycin resistance in *Staphylococci*. *Drug Resist Updates* 1, 135-150.
- Hugh R, Ryschenkow E (1961) *Pseudomonas maltophilia*, an alcaligenes-like species. *Journal of General Microbiology* 26, 123-132.
- Jevitt LA, Smith AJ, Williams PP, Raney PM, McGowan JE, Tenover FC (2003) In vitro activities of daptomycin, linezolid, and quinupristin-dalfopristin against a challenge panel of *Staphylococci* and *Enterococci*, including vancomycin-intermediate *Staphylococcus aureus* and vancomycin-resistant *Enterococcus faecium*. *Microbial Drug Resistance* 9, 389-393.
- Kloos WE, Bannerman TL (1995) *Staphylococcus* and *Micrococcus*. In: Murray PR, Baron EJ, Pfaller MA, Tenover FC, Tenover FC (eds), *Manual of Clinical Microbiology*, 6<sup>th</sup> ed, ASM Press, Washington DC, 282-298.
- Korn-Wendisch F, Kutzner HJ (1992) The family Streptomycetaceae. In: Balows A, Truper HG, Dworkin M, Harder W, Schleifer KH (eds), *The prokaryotes*, Springer-Verlag, New York, 921-995.
- Krueger TS, Clark EA, Nix DE (2001) In vitro susceptibility of *Stenotrophomonas maltophilia* to various antimicrobial combinations. *Diagnostic Microbiology and Infectious Disease* 41, 71-78.
- Lechevalier HA (1989) The actinomycetes III, a practical guide to generic identification of Actinomycetes. *Bergey's Manual of Systematic Bacteriology*, Vol. 4, Williams and Wilkins Company, Baltimore, 2344-2347.
- Levy SB (2002) *The antibiotic paradox: how misuse of antibiotics destroys their curative powers*. 2<sup>nd</sup> (ed), Perseus Books, Boston.
- Levy SB, Marshall B (2004) Antibacterial resistance worldwide: causes, challenges and responses. *Nature Medicine* 10, 122-129.
- Livermore DM (2003) Bacterial resistance: origins, epidemiology and impact. *Clinical Infectious Diseases* 36, 11-23.
- Liu PL, Lau YJ, Hu BS, Shyr JM, Shi ZY, Tsai WS, Lin YH, Tseng CY (1995) Comparison of susceptibility to extended-spectrum  $\beta$ -lactam antibiotics and ciprofloxacin among gram-negative bacilli isolated from intensive care units. *Diagnostic Microbiology and Infectious Disease* 22, 285-291.
- Locci R (1989) *Streptomyces* and related genera. *Bergey's Manual of Systematic Bacteriology*. Vol. 4, Williams and Wilkins Company, Baltimore, 2451-2508.
- Lyczak JB, Cannon CL, Pier GB (2000) Establishment of *Pseudomonas aeruginosa* infection: lessons from a versatile opportunist. *Microbes and Infection* 2, 9, 1051-1060.
- Lyon BR, Skurray R (1987) Antimicrobial resistance of *Staphylococcus aureus*: genetic basis. *Microbiological Reviews* 51, 88-134.
- Madigan MT, Martinko JM, Parker J (1997) Antibiotics: isolation and characterization. In: *Brock Biology of Microorganisms*, 8<sup>th</sup> (ed), Prentice-Hall International Inc. New Jersey, 440-442.
- Meka VG, Gold HS (2004) Antimicrobial resistance to linezolid. *Clinical Infectious Diseases* 39, 1010-1015.
- Miyadoh S (1993) Research on antibiotic screening in Japan over the last decade: A producing microorganisms approach. *Actinomycetologica* 9, 100-106.

- Nathwani D (2005) Tigecycline: clinical evidence and formulary positioning. *International Journal of Antimicrobial Agents* 25, 185-192.
- Nonomura H (1974) Key for classification and identification of 458 species of the *Streptomyces* included in ISP. *Journal of Fermentation Technology* 52, 78-92.
- Obritsch MD, Fish DN, MacLaren R, Jung R (2004) National surveillance of antimicrobial resistance in *Pseudomonas aeruginosa* isolates obtained from Intensive Care Unit Patients from 1993 to 2002. *Antimicrobial Agents and Chemotherapy* 48, 4606-4610.
- Oskay M, Tamer AU, Azeri C (2004) Antibacterial activity of some Actinomycetes isolated from farming soils of Turkey. *African Journal of Biotechnology* 3, 9, 441- 446.
- Palleroni NJ, Bradbury JF (1993) *Stenotrophomonas*, a new bacterial genus for *Xanthomonas maltophilia* (Hugh 1980) Swings et al. 1983. *International Journal of Systematic Bacteriology* 43, 606-609.
- Paterson DL, Ko WC, Von Gottberg A, Mohapatra S, Casellas JM, Goessens H, Mulazimoglu L, Trenholme G, Klugma KP, Bonomo RA, Rice LB, Wagener MM, McCormack JG, Yu VL (2004) International prospective study of *Klebsiella pneumoniae* bacteremia: implications of extended-spectrum beta-lactamase production in nosocomial infections. *Annals of Internal Medicine* 140, 26-32.
- Prauser H (1964) Aptness and application of colour for exact description of colours of *Streptomyces*. *Zeitschrift für allgemeine Mikrobiologie* 4, 95-98.
- Projan S (2002) New (and not so new) antibacterial targets- from where and when will the novel drugs come? *Current Opinion in Pharmacology* 2, 513-522.
- Robin T, Janda JM (1996) Pseudo-, Xantho-, *Stenotrophomonas maltophilia*: an emerging pathogen in search of a genus. *Clinical Microbiology Newsletter* 18, 2, 9- 13.
- Rubin RJ, Harrington CA, Poon A, Dietrich K, Greene JA, Moiduddin A (1999) The economic impact of *Staphylococcus* infection in New York City hospitals. *Emerging Infectious Diseases* 5, 9-17.
- Saiman L, Chen Y, Gabriel PS, Knirsch C (2002) Synergistic activities of macrolide antibiotic against *Pseudomonas aeruginosa*, *Burkholderia cepacia*, *Stenotrophomonas maltophilia*, and *Alcaligenes xylosoxidans* isolated from patients with cystic fibrosis. *Antimicrobial Agents and Chemotherapy* 46, 4, 1105-1107.
- Sahin N, Ugur A, (2003) Investigation of the antimicrobial activity of some *Streptomyces* isolates. *Turkish Journal of Biology* 27, 79-84.
- Shirling EB, Gottlieb D (1966) Methods, classification, identification and description of genera and species. Vol. 2, The Williams and Wilkins Company, Baltimore, 61-292.
- Skaehill P (2000) Management of *Stenotrophomonas maltophilia* infections. *The Consult Pharmacist* 15, 74-76.
- Spizek J, Tichy P (1995) Some aspects of overproduction of secondary metabolites. *Folia Microbiological* 40, 43-50.
- Swings J, Devos M, Van der Mooter M, De Ley J (1983) Transfer of *Pseudomonas maltophilia* Hugh 1981 to the genus *Xanthomonas* as *Xanthomonas maltophilia* (Hugh 1981) comb. nov. *International Journal of Systematic Bacteriology* 33, 409-413.
- Tanaka YT, Mura SO (1993) Agro active compounds of microbial origin. *Annual Review of Microbiology* 47, 57-87.

- Todd JK (1998) Toxic shock syndrome. *Clinical Microbiology Reviews* 1, 432-446.
- Ugur A, Ceylan O (2003) Occurrence of resistance to antibiotics, metals, and plasmids in clinical strains of *Staphylococcus* spp. *Archives of Medical Research* 34, 130-136.
- Urban C, Segal-Maurer S, Rahal JJ (2003) Considerations in control and treatment of nosocomial infections due to multi-drug resistant *Acinetobacter baumannii*. *Clinical Infectious Diseases* 36, 1268-1274.
- Vartivarian S, Anaissie E, Bodey G, Sprigg H, Rolston K (1994) A changing pattern of susceptibility of *Xanthomonas maltophilia* to antimicrobial agents: implications for therapy. *Antimicrobial Agents and Chemotherapy* 38, 624- 627.
- Waksman SA (1961) The Actinomycetes resistant bacteria. *Journal of Arid Environments* 53, 365-371.
- Wenzel RP (2004) The antibiotic pipeline- challenges, costs, and values. *New England Journal of Medicine* 351, 523-526.
- Willett HP (1992) *Staphylococcus*. In: Joklik WK, Willett HP, Amos DB, Wilfert CM (eds), *Zinsser Microbiology*, 20th ed., Norwalk, Appleton and Lange, 401-416.
- Williams ST, Goodfellow M, Alderson G, Wellington EMH, Sneath PHA, Sackin MJ (1983a) Numerical classification of *Streptomyces* and related genera. *Journal of General Microbiology* 129, 1743-1813.
- Williams ST, Goodfellow M, Wellington EMH, Vickers JC, Alderson G, Sneath PHA, Sackin MJ, Mortimer AM (1983b) A probability matrix for identification of *Streptomyces*. *Journal of General Microbiology* 129, 1815-1830.
- Williams ST, Goodfellow M, Alderson G (1989) Genus *Streptomyces* Waksman and Henrici 1943, 339AL. In: Williams ST, Sharpe ME, Holt JG (eds.) *Bergey's Manual of Systematic Bacteriology*, Vol. 4, Williams and Wilkins, Baltimore, 2452-2492.

---

## **Antibiyotik Dirençli Bakterilere Karşı Aktif Antibiyotik Kaynağı Olarak Toprak *Streptomyces*'lerinin İzolasyonu**

### **Ozet**

Bu çalışmanın amacı toprakta yaygın olarak bulunan ve antibiyotik üreticileri olarak bilinen *Streptomyces*'lerin in vitro antimikrobiyal aktiviteleridir. Türkiye, Muğla'daki farklı alanlardan *Streptomyces* izolatları, çoklu antibiyotige dirençli *Staphylococcus aureus* ve *Stenotrophomonas maltophilia*'yi kapsayan on beş mikroorganizma üzerinde inhibitör aktiviteleri değerlendirilmiştir. Test mikroorganizmalarının en az ikisine karşı antimikrobiyal aktivite gösteren on beş *Streptomyces* izolatı geleneksel metodlar ile karakterize edilmiştir. Sonuçlar beş izolatin metisiline dirençli *Staphylococcus aureus* (MRSA)'u kapsayan *S. aureus* suslarına karşı yüksek derecede aktif olduklarını göstermektedir. On iki *Streptomyces* izolatı *Candida albicans*'a karşı antikandidal aktivite göstermiştir. On izolat 30 mm çaptan daha fazla inhibisyon zonlu yüksek aktivitedir. İzolatların çoğu Gram negatif test bakterilerinin gelişimini inhibe etmiştir. Sekiz izolat *S. maltophilia* MU64 üzerinde antibakteriyal aktivite göstermiştir. Bunların ikisinin inhibisyon zonları *S. maltophilia* için 30 mm'den daha yüksektir.

**Anahtar Kelimeler:** Antimikrobiyal aktivite, mikroorganizma, *Streptomyces*.